arXiv:1410.8300v1 [math-ph] 30 Oct 2014

Confluent Heun functions and the Coulomb problem for spin 1/2 particle in Minkowski space

V. Balan*, A.M. Manukyan†, E.M. Ovsiyuk‡, V.M. Red'kov§, O.V. Veko¶

October 31, 2014

Abstract

In the paper, the well-known quantum mechanical problem of a spin 1/2 particle in external Coulomb potential, reduced to a system of two first-order differential equations, is studied from the point of view of possible applications of the Heun function theory to treat this system. It is shown that in addition to the standard way to solve the problem in terms of the confluent hypergeometric functions (proposed in 1928 by G. Darvin and W. Gordon), there are possible several other possibilities which rely on applying the confluent Heun functions. Namely, in the paper there are elaborated two combined possibilities to construct solutions: the first applies when one equation of the pair of relevant functions is expressed trough hypergeometric functions, and another constructed in terms of confluent Heun functions. In this respect, certain relations between the two classes of functions are established. It is shown that both functions of the system may be expressed in terms of confluent Heun functions. All the ways to study this problem lead us to a single energy spectrum, which indicates their correctness.

PACS numbers: 02.30.Gp, 02.30.Hq;
MSC2010: 33E30, 34B30.
Keywords and phrases: Quantum mechanics; Dirac equation; Coulomb problem; confluent Heun functions; confluent hypergeometric functions.

1 Introduction

The general Heun equation is a second order linear differential equation which has four regular singularities and different confluent forms [1, 2]. The general Heun equation and all its confluent forms turn out to be of primary significance in physical applications, for instance in quantum mechanics and field theory on the background of curved space-time models, and in optics – see [3–72]. A more complete and comprehensive list of references can be found on the site of *the Heun Project*[1].

*University Politehnica of Bucharest, Romania, vladimir.balan@upb.ro
†Institute for Physical Research of NAS of Armenia
‡Mosyr State Pedagogical University, Belarus, e.ovsiyuk@mail.ru
§B.I. Stepanov Institute of Physics, NAS of Belarus, redkov@dragon.bas-net.by
¶Kalinkovichi Gymnasium, Belarus,vekoolga@mail.ru

[1]Sofia University, The Heun Project: Heun functions, their generalizations and applications, http://theheunproject.org

In this paper, the well-known quantum mechanical problem of a spin 1/2 particle in external Coulomb potential, reduced to a system of two first-order differential equations, is studied from the point of view of possible applications of the Heun function theory to treat this system. It is shown that in addition to the standard way [73] to solve the problem in terms of the confluent hypergeometric functions (proposed in 1928 by G. Darvin and W. Gordon), there are possible several other possibilities which rely on applying confluent Heun functions. Namely, we shall elaborate two combined possibilities to construct solutions: one for the case when one equation of the pair of relevant functions is expressed trough hypergeometric functions, and another constructed in terms of confluent Heun functions. As well, certain relations between these two classes of functions are established. There exists a possibility to express both the functions of the system in terms of confluent Heun functions. All the ways to study the problem lead us to a unique energy spectrum, which indicates their correctness. In particular physical problems, there exist as well similar possibilities to avoid (with the help of special tricks) the use of Heun functions by restricting to the use of only hypergeometric functions – for instance see in [54, 55, 56, 57].

2 The Coulomb problem: solutions constructed by hypergeometric and partially by Heun functions

In spherical coordinates, a diagonal tetrad has the form [74, 75]

$$\begin{gathered} dS^2 = dt^2 - dr^2 - r^2(d\theta^2 + \sin^2 d\phi^2) \ , \\ e^{\alpha}_{(0)} = (1,0,0,0) \ , \qquad e^{\alpha}_{(3)} = (0,1,0,0) \ , \\ e^{\alpha}_{(1)} = (0,0,r^{-1},0) \ , \qquad e^{\alpha}_{(2)} = (0,0,0,r^{-1}\sin^{-1}\theta) \ . \end{gathered} \tag{1}$$

The Ricci rotation coefficients are

$$\gamma_{ab0} = 0, \ \gamma_{ab3} = 0, \quad \gamma_{ab1} = \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -r^{-1} \\ 0 & 0 & 0 & 0 \\ 0 & +r^{-1} & 0 & 0 \end{array} \right|, \ \gamma_{ab2} = \left| \begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & 0 & \cot\theta r^{-1} & 0 \\ 0 & -\cot\theta r^{-1} & 0 & -r^{-1} \\ 0 & 0 & +r^{-1}\chi & 0 \end{array} \right| .$$

Generally, the covariant Dirac equation[2]

$$\left[i\gamma^c \Big(e^{\alpha}_{(c)} \partial_\alpha + \frac{1}{2} j^{ab} \gamma_{abc} \Big) - m \right] \Psi = 0$$

takes the form

$$\left[i\gamma^0 \frac{\partial}{\partial t} + i\Big(\gamma^3 \frac{\partial}{\partial r} + \frac{\gamma^1 j^{31} + \gamma^2 j^{32}}{r} \Big) + \frac{1}{r}\Sigma_{\theta\phi} - m \right] \Psi = 0, \tag{2}$$

$$\Sigma_{\theta,\phi} = i\gamma^1 \partial_\theta + \gamma^2 \frac{i\partial_\phi \ + i\sigma^{12}}{\sin\theta}. \tag{3}$$

With the help of relations

$$\frac{\gamma^1 j^{31} + \gamma^2 j^{32}}{r} = \frac{\gamma^3}{r}, \quad \Psi = \frac{1}{r}\tilde{\Psi}$$

[2]We further use the notations from [74, 75].

equation (2) is simplified,

$$\left(i\gamma^0\frac{\partial}{\partial t}+i\gamma^3\frac{\partial}{\partial r}+\frac{1}{r}\Sigma_{\theta\phi}-m\right)\tilde{\Psi}=0. \tag{4}$$

In order to diagonalize the operators $i\partial_t, \vec{J}^2, J_3$, one takes the wave function in the form [74]

$$\tilde{\Psi}=e^{-iEt}\left|\begin{array}{c} f_1(\chi)\ D_{-1/2} \\ f_2(\chi)\ D_{+1/2} \\ f_3(\chi)\ D_{-1/2} \\ f_4(\chi)\ D_{+1/2} \end{array}\right|, \tag{5}$$

where the Wigner functions [76] are denoted by $D_\sigma = D^j_{-m.\sigma}(\phi,\theta.0)$. After separating the variables, we get four radial equations (let $\nu = j+1/2$)

$$\begin{array}{ll} Ef_3 - i\dfrac{d}{d\chi}f_3 - i\dfrac{\nu}{\sin\chi}f_4 - mf_1 = 0, & Ef_4 + i\dfrac{d}{d\chi}f_4 + i\dfrac{\nu}{\sin\chi}f_3 - mf_2 = 0, \\ Ef_1 + i\dfrac{d}{d\chi}f_1 + i\dfrac{\nu}{\sin\chi}f_2 - mf_3 = 0, & Ef_2 - i\dfrac{d}{d\chi}f_2 - i\dfrac{\nu}{\sin\chi}f_1 - mf_4 = 0. \end{array} \tag{6}$$

In a spherical tetrad, the space reflection operator is given by [74]

$$\hat{\Pi}_{sph} \ = \left|\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{array}\right| \otimes\ \hat{P}\ .$$

From the eigenvalues equations $\hat{\Pi}_{sph}\ \Psi_{jm} = \ \Pi\ \Psi_{jm}$, we obtain

$$\Pi = \ \delta\ (-1)^{j+1},\ \ \delta = \pm 1\ , \qquad f_4 = \ \delta\ f_1, \qquad f_3 = \ \delta\ f_2\ , \tag{7}$$

which simplifies (6),

$$\left(\frac{d}{dr}+\frac{\nu}{r}\right)\ f + (E+\delta m)\ g = 0, \quad \left(\frac{d}{dr}-\frac{\nu}{r}\right)\ g - (E-\delta m)\ f = 0, \tag{8}$$

where instead of f_1 and f_2, the new variables f and g are used

$$f=\frac{f_1+f_2}{\sqrt{2}}, \qquad g=\frac{f_1-f_2}{i\sqrt{2}}.$$

For definiteness, let us consider the case when $\delta = 1$,

$$\left(\frac{d}{dr}+\frac{\nu}{r}\right)f + (E+m)g = 0, \quad \left(\frac{d}{dr}-\frac{\nu}{r}\right)g - (E-m)f = 0; \tag{9}$$

by performing the replacement $m \rightsquigarrow -m$ we obtain the equations for $\delta = -1$.

The presence of the external Coulomb field is taken into account in (9) by the formal change $\epsilon \rightsquigarrow \epsilon + \frac{e}{r}$. Thus, the quantum Coulomb problem for a Dirac particle is described by the following radial system

$$\left(\frac{d}{dr}+\frac{\nu}{r}\right)f + \left(E+\frac{e}{r}+m\right)g = 0, \quad \left(\frac{d}{dr}-\frac{\nu}{r}\right)g - \left(E+\frac{e}{r}-m\right)f = 0. \tag{10}$$

Let us perform a linear transformation over the functions $f(r)$ and $g(r)$[3]

$$f(r) = aF(r) + cG(r), \quad g(r) = dF(r) + bG(r)\ ,$$
$$F(r) = bf(r) - cg(r), \quad G(r) = -df(r) + ag(r). \tag{11}$$

Let us combine the equations (10) as follows: the first equation is multiplied by $+b$, the second by $-c$, and then sum the results; analogously, we add the first equation multiplied by $-d$ with the second multiplied by $+a$. Thus we arrive at

$$\left[\ \frac{d}{dr} - b'a + c'd + \frac{\nu}{r}\,(ba+cd) + \left(E + \frac{e}{r} + m\right) bd + \left(E + \frac{e}{r} - m\right) ca\ \right] F$$
$$= \left[\ b'c - bc' - \frac{\nu}{r} 2bc - \left(E + \frac{e}{r} + m\right) b^2 - \left(E + \frac{e}{r} - m\right) c^2\ \right] G\ ,$$
$$\left[\ \frac{d}{dr} + d'c - a'b - \frac{\nu}{r}\,(dc+ab) - \left(E + \frac{e}{r} + m\right) bd - \left(E + \frac{e}{r} - m\right) ca\ \right] G$$
$$= \left[\ -d'a + da' + \frac{\nu}{r} 2ad + \left(E + \frac{e}{r} + m\right) d^2 + \left(E + \frac{e}{r} - m\right) a^2\ \right] F\ . \tag{12}$$

For simplicity, let us assume that the transformation (11) does not depend on r, and let it be orthogonal:

$$S = \left|\begin{array}{cc} a & c \\ d & b \end{array}\right| = \left|\begin{array}{cc} \cos A/2 & \sin A/2 \\ -\sin A/2 & \cos A/2 \end{array}\right|, \tag{13}$$

which simplifies the equations (12)

$$\left(\frac{d}{dr} + \frac{\nu}{r}\cos A - m\sin A\right) F = \left(-\frac{\nu}{r}\sin A - \frac{e}{r} - E - m\cos A\right) G,$$
$$\left(\frac{d}{dr} - \frac{\nu}{r}\cos A + m\sin A\right) G = \left(-\frac{\nu}{r}\sin A + \frac{e}{r} + E - m\cos A\right) F. \tag{14}$$

There exist four possibilities (only two of them are different in fact):

1)
$$-\frac{\nu}{r}\sin A + \frac{e}{r} = 0\ , \qquad \sin A = \frac{e}{\nu}\ , \qquad \cos A = \sqrt{1 - e^2/\nu^2}\ ,$$
$$\cos\frac{A}{2} = \sqrt{\frac{\nu + \sqrt{\nu^2 - e^2}}{2\nu}}\ , \qquad \sin\frac{A}{2} = \sqrt{\frac{\nu - \sqrt{\nu^2 - e^2}}{2\nu}}\ ;$$

1′)
$$-\frac{\nu}{r}\sin A - \frac{e}{r} = 0\ , \qquad \sin A = -\frac{e}{\nu}, \quad \cos A = \sqrt{1 - e^2/\nu^2},$$
$$\cos\frac{A}{2} = \sqrt{\frac{\nu - \sqrt{\nu^2 - e^2}}{2\nu}}, \quad \sin\frac{A}{2} = \sqrt{\frac{\nu + \sqrt{\nu^2 - e^2}}{2\nu}}\ ; \tag{15}$$

2)
$$E - m\cos A = 0, \quad \cos A = +\frac{E}{m}, \quad \sin A = \sqrt{1 - E^2/m^2},$$

[3]The coefficients of this transformation may depend on the radial variable; let its determinant obey the identity $a(r)b(r) - c(r)d(r) = 1$.

$$\cos\frac{A}{2}=\sqrt{\frac{m+E}{2m}},\qquad \sin\frac{A}{2}=\sqrt{\frac{m-E}{2m}}\;;$$

$2'$)

$$-E-m\cos A=0,\qquad \cos A=-\frac{E}{m},\quad \sin A=\sqrt{1-E^2/m^2},$$
$$\cos\frac{A}{2}=\sqrt{\frac{m-E}{2m}},\quad \sin\frac{A}{2}=\sqrt{\frac{m+E}{2m}}\,. \tag{16}$$

First, consider the case 1). Equations (14) take the form

$$\left(\frac{d}{dr}+\frac{\nu}{r}\cos A-m\sin A\right)F=\left(-\frac{2e}{r}-E-m\cos A\right)G,$$
$$\left(\frac{d}{dr}-\frac{\nu}{r}\cos A+m\sin A\right)G=(E-m\cos A)F. \tag{17}$$

After eliminating the function F, we get a second order equation for G

$$\left(\frac{d}{dr}+\frac{\nu}{r}\cos A-m\sin A\right)\left(\frac{d}{dr}-\frac{\nu}{r}\cos A+m\sin A\right)G$$
$$=(E-m\cos A)\left(-\frac{2e}{r}-E-m\cos A\right)G, \tag{18}$$

or, differently,

$$\left(\frac{d^2}{dr^2}+E^2-m^2+\frac{\nu\cos A-\nu^2\cos^2 A}{r^2}+\frac{2eE-2em\cos A+2m\nu\sin A\cos A}{r}\right)G=0.$$

Having in mind the identity $\sin A=e/\nu$, the last equation reduces to

$$\left(\frac{d^2}{dr^2}+E^2-m^2+\frac{\nu\cos A-\nu^2\cos^2 A}{r^2}+\frac{2eE}{r}\right)G=0. \tag{19}$$

After changing the variable, $x=2\sqrt{m^2-E^2}\,r$, it reads

$$\frac{d^2G}{dx^2}+\left(-\frac{1}{4}-\frac{\nu\cos A\,(\nu\cos A-1)}{x^2}+\frac{eE}{\sqrt{m^2-E^2}\,x}\right)G=0\,. \tag{20}$$

With the use of the substitution $G(x)=x^a e^{bx}\bar{G}(x)$ for $\bar{G}$, we get

$$x\,\frac{d^2\bar{G}}{dx^2}+(2\,a+2\,b\,x)\,\frac{d\bar{G}}{dx}$$
$$+\left[(b^2-\frac{1}{4})\,x+\frac{a^2-a-\nu\cos A\;(\nu\cos A-1)}{x}+2ab+\frac{eE}{\sqrt{m^2-E^2}}\right]\bar{G}=0\,.$$

When

$$a=+\nu\,\cos A=\sqrt{\nu^2-e^2}\,,\qquad b=-\frac{1}{2}\,,$$

this equation for $\bar{G}$ becomes simpler,

$$x\,\frac{d^2\bar{G}}{dx^2}+(2\,a-\,x)\,\frac{d\bar{G}}{dx}-\left(a-\frac{eE}{\sqrt{m^2-E^2}}\right)\varphi=0\,,$$

which is a confluent hypergeometric equation

$$x\;{}_1F_1''(x)+(\gamma-x){}_1F_1'(x)-\alpha\;{}_1F_1(x)=0\;,\qquad \alpha=a-\frac{eE}{\sqrt{m^2-E^2}}\,,\qquad \gamma=2a\,.$$

To polynomials there corresponds the known restriction $\alpha=-n$, $n=0,1,2,...$, which gives the known energy quantization rule

$$a-\frac{eE}{\sqrt{m^2-E^2}}=-n\qquad\Longrightarrow\qquad E=\frac{m}{\sqrt{1+e^2/(n+\sqrt{\nu^2-e^2})^2}}\;. \tag{21}$$

In turn, from (17) it follows a second order equation for F,

$$\left(\frac{d}{dr}-\frac{\nu}{r}\;\cos A+m\;\sin A\right)\frac{r}{2e+(E+m\cos A)r}\left(\frac{d}{dr}+\frac{\nu}{r}\;\cos A-m\;\sin A\right)F$$

$$=(-E+m\cos A)F\;, \tag{22}$$

or

$$\left[\Big(\frac{d}{dr}-\frac{\nu}{r}\;\cos A+m\;\sin A\Big)\Big(\frac{d}{dr}+\frac{\nu}{r}\;\cos A-m\;\sin A\Big)\right.$$
$$\left.+\frac{d}{dr}\ln\left(\frac{r}{2e+(E+m\cos A)r}\right)\Big(\frac{d}{dr}+\frac{\nu}{r}\;\cos A-m\;\sin A\Big)\right]F$$
$$=\Big(\frac{2e}{r}+E+m\cos A\Big)\;\Big(-E+m\cos A\Big)\;F\;,$$

and further,

$$\left[\frac{d^2}{dr^2}+\left(\frac{1}{r}-\frac{E+m\cos A}{2e+(E+m\cos A)r}\right)\frac{d}{dr}\right.$$
$$+\frac{\nu}{r^2}\;\cos A-\frac{m\;\sin A}{r}-\left(\frac{1}{r}-\frac{2e}{r[2e+(E+m\cos A)r]}\right)\Big(\frac{\nu}{r}\;\cos A-m\;\sin A\Big)$$
$$\left.+E^2-m^2-\frac{\nu\cos A}{r^2}+\frac{e^2-\nu^2}{r^2}+\frac{2eE}{r}\right]F=0\;.$$

Finally, we obtain

$$\left[\frac{d^2}{dr^2}+\Big(\frac{1}{r}-\frac{E+m\cos A}{2e+(E+m\cos A)r}\Big)\frac{d}{dr}\right.$$
$$\left.+E^2-m^2+\frac{2eE}{r}+\frac{e^2-\nu^2}{r^2}-\frac{2em\sin A+\nu\cos A(E+m\cos A)}{r\;[2e+(E+m\cos A)r]}\right]F=0\;.$$

Let us introduce special designation for the additional singular point

$$-\frac{2e}{E+m\cos A}=R\;;$$

then we obtain

$$\left[\frac{d^2}{dr^2}+\left(\frac{1}{r}-\frac{1}{r-R}\right)\frac{d}{dr}+E^2-m^2+\frac{2eE}{r}+\frac{e^2-\nu^2}{r^2}+\frac{mR\,\sin A-\nu\cos A}{r\,(r-R)}\right]F=0. \tag{23}$$

After changing the variable $y=r/R$, it reads

$$\frac{d^2F}{dy^2}+\left(\frac{1}{y}-\frac{1}{y-1}\right)\frac{dF}{dy}+\left((E^2-m^2)\,R^2-\frac{\nu^2-e^2}{y^2}\right.$$
$$\left.+\frac{-\nu\cos A+mR\,\sin A}{y-1}+\frac{2eRE-mR\sin A+\nu\cos A}{y}\right)F=0\,. \tag{24}$$

Let us search solutions in the form $F=y^a\,e^{by}\,\bar{F}(y)$; the function $\bar{F}$ obeys

$$\frac{d^2\bar{F}}{dy^2}+\left(\frac{2a+1}{y}+2b-\frac{1}{y-1}\right)\frac{d\bar{F}}{dy}$$
$$+\left[b^2+(E^2-m^2)\,R^2+\frac{a^2-\nu^2+e^2}{y^2}-\frac{a+b+\nu\cos A-mR\,\sin A}{y-1}\right.$$
$$\left.+\frac{a+b+2ab+R(2eE-m\sin A)+\nu\cos A}{y}\right]\bar{F}=0\,.$$

With a,b taken according to[4]:

$$a=\underline{+\sqrt{\nu^2-e^2}}\,,\;\;-\sqrt{\nu^2-e^2}\,,$$
$$b=+\sqrt{m^2-E^2}\,R\,,\underline{-\sqrt{m^2-E^2}\,R} \tag{25}$$

the above equation becomes simpler

$$\frac{d^2\bar{F}}{dy^2}+\left(2b+\frac{2a+1}{y}-\frac{1}{y-1}\right)\frac{d\bar{F}}{dy}$$
$$+\left(\frac{a+b+2ab+2eRE-mR\sin A+\nu\cos A}{y}\right.$$
$$\left.-\frac{a+b+\nu\cos A-mR\,\sin A}{y-1}\right)\bar{F}=0\,. \tag{26}$$

This can be easily recognized as a confluent Heun equation for $H(\alpha,\beta,\gamma,\delta,\eta,z)$,

$$H''+\left(\alpha+\frac{1+\beta}{z}+\frac{1+\gamma}{z-1}\right)H'+\left(\frac{1}{2}\,\frac{\alpha+\alpha\beta-\beta-\beta\gamma-\gamma-2\eta}{z}\right.$$
$$\left.+\frac{1}{2}\,\frac{\alpha+\alpha\gamma+\beta+\beta\gamma+\gamma+2\delta+2\eta}{z-1}\right)H=0 \tag{27}$$

with the parameters

$$\alpha=2b\,,\qquad\beta=2a\,,\qquad\gamma=-2\,,$$
$$\delta=2\,eER\,,\qquad\eta=1+mR\,\sin A-2\,eER-\nu\cos A\,. \tag{28}$$

[4]We shall use below underlined values.

Let us use the known condition [2] (one of two conditions needed to get polynomials):

$$\delta = -\left(n + \frac{\beta+\gamma+2}{2}\right)\alpha\,, \qquad n = 0, 1, 2, \ldots\,, \tag{29}$$

it results the energy quantization rule

$$a = +\sqrt{\nu^2 - e^2}\,, \qquad b = -\sqrt{m^2 - E^2}\,R\,,$$
$$eER = (n + \sqrt{\nu^2 - e^2})\,\sqrt{m^2 - E^2}\;R\,,$$

so we infer

$$E = \frac{m}{\sqrt{1 + e^2/(n + \sqrt{\nu^2 - e^2})^2}}\,, \tag{30}$$

which coincides with the known formula for energy levels.

It should be emphasized that as follows from (17), the function F (being constructed in terms of the confluent Heun functions) can be related with the function G (which is determined in terms of confluent hypergeometric functions) by means of the following differential operators:

$$G = \left(-\frac{2e}{r} - E - m\cos A\right)^{-1}\left(\frac{d}{dr} + \frac{\nu}{r}\cos A - m\sin A\right)F, \tag{31}$$

and

$$F = \frac{1}{(E - m\cos A)}\left(\frac{d}{dr} - \frac{\nu}{r}\cos A + m\sin A\right)G. \tag{32}$$

They can be rewritten in the form

$$G(r) = \frac{1}{2e}\frac{y}{y-1}\left(\frac{d}{dy} + \frac{\nu}{y}\cos A - mR\sin A\right)F(y), \tag{33}$$

and

$$F(y) = \frac{r}{2(Er+e)}\left(\frac{d}{dr} - \frac{\nu}{r}\cos A + m\sin A\right)G(r). \tag{34}$$

Let us consider the case 2) – see (16). The equations (14) take the form

$$\left(\frac{d}{dr} + \frac{\nu}{r}\;\cos A - m\;\sin A\right)F = \left(-\frac{\nu\sin A + e}{r} - 2m\cos A\right)G\,,$$
$$\left(\frac{d}{dr} - \frac{\nu}{r}\;\cos A + m\;\sin A\right)G = \frac{e - \nu\sin A}{r}\;F\,. \tag{35}$$

One can obtain a second order equation for $G(r)$:

$$\Big[\Big(\frac{d}{dr} + \frac{\nu}{r}\;\cos A - m\;\sin A\Big)\;r\Big(\frac{d}{dr} - \frac{\nu}{r}\;\cos A + m\;\sin A\Big)$$
$$+(e - \nu\sin A)\Big(\frac{e + \nu\sin A}{r} + 2m\cos A\Big)\Big]\;G = 0\,, \tag{36}$$

from where it follows

$$\left[\frac{d}{dr}-\frac{\nu}{r}\,\cos A+m\,\sin A+r\left(\frac{d^2}{dr^2}+\frac{\nu\cos A}{r^2}\,-\left(\frac{\nu}{r}\,\cos A-m\,\sin A\right)^2\right)\right.$$

$$\left.+(e-\nu\sin A)\left(\frac{e+\nu\sin A}{r}+2m\cos A\right)\right]G=0\,,$$

or

$$\left(\frac{d^2}{dr^2}+\frac{1}{r}\frac{d}{dr}-m^2\,\sin^2 A+\frac{e^2-\nu^2}{r^2}+\frac{2me\cos A}{r}+\frac{m\sin A}{r}\right)G=0\,.$$

Further, taking into account the identity $\cos A=E/m$, we get

$$\left(\frac{d^2}{dr^2}+\frac{1}{r}\frac{d}{dr}+E^2-m^2+\frac{e^2-\nu^2}{r^2}+\frac{2eE}{r}+\frac{\sqrt{m^2-E^2}}{r}\right)G=0\,. \tag{37}$$

By making the change of variables $x=2\sqrt{m^2-E^2}\,r$, we get

$$\frac{d^2G}{dx^2}+\frac{1}{x}\,\frac{dG}{dx}+\left(-\frac{1}{4}-\frac{\nu^2-e^2}{x^2}+\frac{1}{2}\,\frac{m^2-E^2+2Ee\sqrt{m^2-E^2}}{(m^2-E^2)\,x}\right)G=0\,.$$

Let $G(x)=x^a e^{bx}\bar{G}(x)$; the function $\bar{G}$ satisfies

$$x\,\frac{d^2\bar{G}}{dx^2}+(2\,a+1+2\,b\,x)\,\frac{d\bar{G}}{dx}$$

$$+\left[\left(b^2-\frac{1}{4}\right)x+\frac{a^2-\nu^2+e^2}{x}+2ab+b+\frac{1}{2}\,\frac{m^2-E^2+2Ee\sqrt{m^2-E^2}}{m^2-E^2}\right]\bar{G}=0\,.$$

When

$$a=\sqrt{\nu^2-e^2}\,,\qquad b=-\frac{1}{2}\,;$$

we get

$$x\,\frac{d^2\varphi}{dx^2}+(2\,a+1-x)\,\frac{d\varphi}{dx}-\left(a-\,\frac{Ee}{\sqrt{m^2-E^2}}\right)\varphi=0\,,$$

which is a confluent hypergeometric equation

$$x\;{}_1F_1''+(\gamma-x){}_1F_1'-\alpha\;{}_1F_1=0\,,\qquad \alpha=a-\,\frac{Ee}{\sqrt{m^2-E^2}}\,,\qquad \gamma=2a+1\,.$$

The solutions become polynomials if $\alpha=-n,\ n=0,1,2,...$; this provides us with the energy spectrum

$$E=\frac{m}{\sqrt{1+e^2/(n+\sqrt{\nu^2-e^2})^2}}\,. \tag{38}$$

In turn, from (35) it follows a second order equation for $F(r)$

$$\left(\frac{d}{dr}-\frac{\nu}{r}\,\cos A+m\,\sin A\right)\frac{r}{\nu\sin A+e+2m\cos A\,r}\left(\frac{d}{dr}+\frac{\nu}{r}\,\cos A-m\,\sin A\right)F$$

$$= \frac{\nu \sin A - e}{r}\, F\ ,$$

that is

$$\Big[\Big(\frac{d}{dr} - \frac{\nu}{r}\,\cos A + m\,\sin A\Big)\,\Big(\frac{d}{dr} + \frac{\nu}{r}\,\cos A - m\,\sin A\Big)$$
$$+\left(\frac{d}{dr}\ln\frac{r}{\nu\sin A + e + 2m\cos A\, r}\right)\Big(\frac{d}{dr} + \frac{\nu}{r}\,\cos A - m\,\sin A\Big)$$
$$+\frac{e + \nu\sin A + 2m\cos A\, r}{r}\,\frac{e - \nu\sin A}{r}\Big]\, F = 0\ .$$

After simple transformations, we obtain

$$\Big[\frac{d^2}{dr^2} + \left(\frac{1}{r} - \frac{2m\cos A}{\nu\sin A + e + 2m\cos A\, r}\right)\frac{d}{dr}$$
$$-\frac{2m\cos A}{\nu\sin A + e + 2m\cos A\, r}\Big(\frac{\nu}{r}\,\cos A - m\,\sin A\Big)$$
$$+E^2 - m^2 + \frac{e^2 - \nu^2}{r^2} + \frac{2eE - m\sin A}{r}\Big]\, F = 0\ .$$

With the following special notation for the additional singular point

$$D = -\frac{e + \nu\sin A}{2m\cos A}\ , \tag{39}$$

we get the shorter form

$$\Big[\frac{d^2}{dr^2} + \left(\frac{1}{r} - \frac{1}{r-D}\right)\frac{d}{dr} + \frac{m\sin A}{r-D} - \frac{\nu\cos A}{D}\Big(\frac{1}{r-D} - \frac{1}{r}\Big)$$
$$+E^2 - m^2 + \frac{e^2 - \nu^2}{r^2} + \frac{2eE - m\sin A}{r}\Big]\, F = 0\ . \tag{40}$$

Relative to the variable $y = r/D$, this looks simpler

$$\frac{d^2F}{dy^2} + \left[\frac{1}{y} - \frac{1}{y-1}\right]\,\frac{dF}{dy} + \Big[(E^2 - m^2)\, D^2 - \frac{\nu^2 - e^2}{y^2}$$
$$+\frac{-\nu\cos A + mD\,\sin A}{y-1} + \frac{D(2eE - m\sin A) + \nu\cos A}{y}\Big]\, F = 0\,. \tag{41}$$

Let $F = y^a\, e^{by}\, \bar{F}(y)$; the function $\bar{F}$ satisfies

$$\frac{d^2\bar{F}}{dy^2} + \left(\frac{2a+1}{y} + 2b - \frac{1}{y-1}\right)\,\frac{d\bar{F}}{dy}$$
$$+\Big[b^2 + (E^2 - m^2)\, D^2 + \frac{a^2 - \nu^2 + e^2}{y^2} - \frac{a + b + \nu\cos A - mD\,\sin A}{y-1}$$
$$+\frac{a + b + 2ab + D(2eE - m\sin A) + \nu\cos A}{y}\Big]\,\bar{F} = 0\,.$$

When a and b are[5]:

$$a = \underline{+\sqrt{\nu^2 - e^2}}\,, \;\; -\sqrt{\nu^2 - e^2}\,,$$
$$b = +\sqrt{m^2 - E^2}\,D\,, \;\; \underline{-\sqrt{m^2 - E^2}\,D}\,, \tag{42}$$

this reads

$$\frac{d^2\bar{F}}{dy^2} + \left(2b + \frac{2a+1}{y} - \frac{1}{y-1}\right)\frac{d\bar{F}}{dy}$$
$$+ \left(\frac{a + b + 2ab + D(2eE - m\sin A) + \nu\cos A}{y}\right.$$
$$\left. - \frac{a + b + \nu\cos A - mD\,\sin A}{y-1}\right)\bar{F} = 0\,, \tag{43}$$

which is a confluent Heun equation for $H(\alpha, \beta, \gamma, \delta, \eta, y)$

$$H'' + \left(\alpha + \frac{1+\beta}{y} + \frac{1+\gamma}{y-1}\right)H'$$
$$+ \left(\frac{1}{2}\,\frac{\alpha + \alpha\beta - \beta - \beta\gamma - \gamma - 2\eta}{y} + \frac{1}{2}\,\frac{\alpha + \alpha\gamma + \beta + \beta\gamma + \gamma + 2\delta + 2\eta}{y-1}\right)G = 0\,, \tag{44}$$

with the parameters

$$\alpha = 2b\,, \qquad \beta = 2a\,, \qquad \gamma = -2\,,$$
$$\delta = 2\,eED\,, \qquad \eta = 1 + mD\,\sin A - 2\,eED - \nu\cos A\,. \tag{45}$$

By imposing one of the two conditions for polynomial solutions [2]

$$\delta = -\left(n + \frac{\beta + \gamma + 2}{2}\right)\alpha\,, \qquad n = 0, 1, 2, \ldots \tag{46}$$

we produce the energy quantization rule

$$a = +\sqrt{\nu^2 - e^2}\,, \qquad b = -\sqrt{m^2 - E^2}\,D\,,$$
$$eED = (n + \sqrt{\nu^2 - e^2})\,\sqrt{m^2 - E^2}\;D\,,$$

whence it follows

$$E = \frac{m}{\sqrt{1 + e^2/(n + \sqrt{\nu^2 - e^2})^2}}\,, \tag{47}$$

which coincides with the known exact result.

It should be emphasized that confluent Heun equations from the cases 1) and 2) formally coincide; however all the parameters are in fact different:

$$1) \qquad \alpha = 2b\,, \qquad \beta = 2a\,, \qquad \gamma = -2\,,$$
$$\delta = 2\,eER\,, \qquad \eta = 1 + mR\,\sin A - 2\,eER - \nu\cos A\,, \tag{48}$$
$$2) \qquad \alpha = 2b\,, \qquad \beta = 2a\,, \qquad \gamma = -2\,,$$
$$\delta = 2\,eED\,, \qquad \eta = 1 + mD\,\sin A - 2\,eED - \nu\cos A\,, \tag{49}$$

[5]We shall use below underlined values.

where

$$1)\quad R=-\frac{2e}{E+m\cos A}\,,\qquad \sin A=\frac{e}{\nu}\,,\qquad \cos A=\sqrt{1-\frac{e^2}{\nu^2}}\,; \tag{50}$$

$$2)\quad D=-\frac{e+\nu\sin A}{2E}\,,\qquad \cos A=\frac{E}{m}\,,\qquad \sin A=\sqrt{1-\frac{E^2}{m^2}}\,. \tag{51}$$

3 Standard treatment of the Coulomb problem

It should be emphasized that the both proposed treatments of the Coulomb problem for Dirac equation differ from the well known one[6]. Let us recall this standard approach. To this end, in the radial system (10)

$$\left(\frac{d}{dr}+\frac{\nu}{r}\right)f+\left(E+\frac{e}{r}+m\right)g=0,\quad \left(\frac{d}{dr}-\frac{\nu}{r}\right)g-\left(E+\frac{e}{r}-m\right)f=0\,. \tag{52}$$

one should introduce the new functions

$$f=\sqrt{m+E}\,(F_1+F_2)\,,\quad g=\sqrt{m-E}\,(F_1-F_2)\,; \tag{53}$$

this infers

$$\left(\frac{d}{dr}+\frac{\nu}{r}\right)(F_1+F_2)+\left(E+\frac{e}{r}+m\right)\frac{\sqrt{m-E}}{\sqrt{m+E}}\,(F_1-F_2)=0,$$
$$\left(\frac{d}{dr}-\frac{\nu}{r}\right)(F_1-F_2)-\left(E+\frac{e}{r}-m\right)\frac{\sqrt{m+E}}{\sqrt{m-E}}\,(F_1+F_2)=0.$$

or,

$$r\left(\frac{d}{dr}+\frac{\nu}{r}\right)(F_1+F_2)+r\,\sqrt{m^2-E^2}(F_1-F_2)+e\frac{\sqrt{m-E}}{\sqrt{m+E}}\,(F_1-F_2)=0,$$
$$r\left(\frac{d}{dr}-\frac{\nu}{r}\right)(F_1-F_2)+r\,\sqrt{m^2-E^2}(F_1+F_2)-e\frac{\sqrt{m+E}}{\sqrt{m-E}}\,(F_1+F_2)=0.$$

By summing and subtracting the equations, we obtain

$$r\frac{d}{dr}F_1+\nu F_2+r\sqrt{m^2-E^2}F_1-\frac{eE}{\sqrt{m^2-E^2}}F_1-\frac{em}{\sqrt{m^2-E^2}}F_2=0$$

$$r\frac{d}{dr}F_2+\nu F_1-r\sqrt{m^2-E^2}F_2+\frac{em}{\sqrt{m^2-E^2}}F_1+\frac{eE}{\sqrt{m^2-E^2}}F_2=0$$

In the variables

$$\lambda=\sqrt{m^2-E^2},\quad x=\lambda r,\quad \frac{em}{\lambda}=\mu,\quad \frac{eE}{\lambda}=\epsilon, \tag{54}$$

these equations read shorter

$$\left(x\frac{d}{dx}+x-\epsilon\right)F_1+(\nu-\mu)\,F_2=0,\quad \left(x\frac{d}{dx}-x+\epsilon\right)F_2+(\nu+\mu)\,F_1=0\,. \tag{55}$$

[6]This was firstly given by G. Darvin and W. Gordon (1928)) – see in [73].

The system (55) can be solved in hypergeometric functions [73].

To detail this point, translate eqs. (55) to a new variable $y = 2x$:

$$\left(y\frac{d}{dy} + \frac{y}{2} - \epsilon\right) F_1 + (\nu - \mu)\, F_2 = 0, \quad \left(y\frac{d}{dy} - \frac{y}{2} + \epsilon\right) F_2 + (\nu + \mu)\, F_1 = 0\,. \tag{56}$$

From whence it follows second order differential equations for F_1 and F_2:

$$\left(y\frac{d^2}{dy^2} + \frac{d}{dy} + \epsilon + \frac{1}{2} - \frac{y}{4} + \frac{\mu^2 - \nu^2 - \epsilon^2}{y}\right) F_1 = 0\,, \tag{57}$$

$$\left(y\frac{d^2}{dy^2} + \frac{d}{dy} + \epsilon - \frac{1}{2} - \frac{y}{4} + \frac{\mu^2 - \nu^2 - \epsilon^2}{y}\right) F_2 = 0\,. \tag{58}$$

Let us study eq. (57). With the substitution $F_1 = y^A e^{By} f_1$ it gives

$$\left[y\frac{d^2}{dy^2} + (2A + 1 + 2By)\ \frac{d}{dy} + \epsilon + \frac{1}{2} + B\ (1 + 2A) - y\left(\frac{1}{4} - B^2\right) + \frac{A^2 + \mu^2 - \nu^2 - \epsilon^2}{y}\right] f_1 = 0\,. \tag{59}$$

The choice

$$A = +\sqrt{\epsilon^2 - \mu^2 + \nu^2}\,, \qquad B = -\frac{1}{2}$$

simplifies eq. (59)to that of confluent hypergeometric type

$$\left[y\frac{d^2}{dy^2} + (2A + 1 - y)\ \frac{d}{dy} + \epsilon - A\right] f_1 = 0 \tag{60}$$

with parameters

$$\alpha_1 = A - \epsilon\,, \qquad \gamma_1 = 2A + 1\,.$$

Imposing the polynomial condition $\alpha_1 = -n_1$ we obtain the quantization rule for ϵ:

$$-\,n_1 = -\epsilon + \sqrt{\epsilon^2 - \mu^2 + \nu^2}\,, \tag{61}$$

Allowing for the above definitions

$$\lambda = \sqrt{m^2 - E^2}, \quad \frac{em}{\lambda} = \mu, \quad \frac{eE}{\lambda} = \epsilon\,;$$

we get

$$\sqrt{\epsilon^2 - \mu^2 + \nu^2} = \sqrt{\nu^2 - e^2}\,, \quad A = +\sqrt{\nu^2 - e^2}\,, \tag{62}$$

and further from (66) we derive a formula for energy levels

$$\frac{eE}{\sqrt{m^2 - E^2}} = \sqrt{\nu^2 - e^2} + n_2 \equiv N_1 \quad \Longrightarrow \quad E = \frac{m}{\sqrt{1 + e^2/N_1^2}}\,, \tag{63}$$

which coincides with (47).

Now, let us consider second equation (58). With the use of the substitution $F_2 = y^a e^{by} f_2$:

$$\left[y\frac{d^2}{dy^2} + (2a+1+2by)\,\frac{d}{dy} + \epsilon - \frac{1}{2} + b\,(1+2a) - y\left(\frac{1}{4} - b^2\right) + \frac{a^2+\mu^2-\nu^2-\epsilon^2}{y} \right] f_2 = 0\,. \quad (64)$$

at

$$a = \pm\sqrt{\epsilon^2 - \mu^2 + \nu^2}\,, \qquad b = -\frac{1}{2}$$

we obtain an equation of the confluent hypergeometric type

$$\left[y\frac{d^2}{dy^2} + (2a+1-y)\,\frac{d}{dy} + \epsilon - a - 1 \right] f_2 = 0 \quad (65)$$

with parameters

$$\alpha_2 = a + 1 - \epsilon\,, \qquad \gamma_2 = 2a + 1\,.$$

Imposing the polynomial restriction $\alpha_2 = -n_2$ we get the quantization rule for ϵ:

$$-\,n_2 = -\epsilon + 1 + \sqrt{\epsilon^2 - \mu^2 + \nu^2}\,; \quad (66)$$

so we arrive at the formula for energy levels

$$\frac{eE}{\sqrt{m^2 - E^2}} = 1 + \sqrt{\nu^2 - e^2} + n_2 \equiv N_2 \quad \Longrightarrow \quad E = \frac{m}{\sqrt{1 + e^2/N_2^2}}\,. \quad (67)$$

Let us find a relative coefficient between two functions:

$$\left(y\frac{d}{dy} + \frac{y}{2} - \epsilon \right) F_1 + (\nu - \mu)\, F_2 = 0,$$
$$\left(y\frac{d}{dy} - \frac{y}{2} + \epsilon \right) F_2 + (\nu + \mu)\, F_1 = 0\,. \quad (68)$$

$$F_1 = C_1 y^A e^{-y/2} F(-n_1, \gamma, y), \quad F_2 = C_2 y^A e^{-y/2} F(-n_2, \gamma, y),,$$
$$A = +\sqrt{\epsilon^2 - \mu^2 + \nu^2}\,, \quad \gamma = 2A = 1, \quad -n_2 = -n_1 + 1\,. \quad (69)$$

We substitute the expressions for the functions F_1, F_2 in the first-order equation (68)

$$y\frac{d}{dy}\left[C_1\, y^A e^{-y/2} F(-n_1, \gamma, y) \right] + \frac{y}{2}\, C_1\, y^A e^{-y/2} F(-n_1, \gamma, y) -$$
$$-\epsilon\, C_1\, y^A e^{-y/2} F(-n_1, \gamma, y) + (\nu - \mu)\, C_2\, y^A e^{-y/2} F(-n_2, \gamma, y) = 0,$$

or

$$y\, C_1\, \frac{d}{dy}\, F(-n_1, \gamma, y) + C_1\,(A - \epsilon)\, F(-n_1, \gamma, y) +$$
$$+ (\nu - \mu)\, C_2\, F(-n_1 + 1, \gamma, y) = 0.$$

We apply the rule of differentiation of the confluent hypergeometric function

$$\frac{d}{dy}\,F(-n_1,\gamma,y)=-\frac{n_1}{y}\,F(-n_1+1,\gamma,y)+\frac{n_1}{y}\,F(-n_1,\gamma,y)\,,$$

as a result we obtain

$$-\,C_1\,n_1\,F(-n_1+1,\gamma,y)+C_1\,n_1\,F(-n_1,\gamma,y)+C_1\,(A-\epsilon)\,F(-n_1,\gamma,y)+\\ +\,(\nu-\mu)\,C_2\,F(-n_1+1,\gamma,y)=0.$$

Taking into account $-n_1=A-\epsilon$, we obtain

$$\frac{C_1}{C_2}=\frac{\nu-\mu}{n_1}=-\frac{\nu-\mu}{A-\epsilon}. \tag{70}$$

Now, we substitute the expressions for the functions F_1, F_2 in the first-order equation (69)

$$y\frac{d}{dy}\left[C_2\,y^A e^{-y/2}F(-n_2,\gamma,y)\right]-\frac{y}{2}\,C_2\,y^A e^{-y/2}F(-n_2,\gamma,y)+\\ +\epsilon\,C_2\,y^A e^{-y/2}F(-n_2,\gamma,y)+(\nu+\mu)\,C_1\,y^A e^{-y/2}F(-n_1,\gamma,y)=0\,,$$

or

$$y\,C_2\,\frac{d}{dy}F(-n_1+1,\gamma,y)+C_2\,(A+\epsilon)\,F(-n_1+1,\gamma,y)-\\ -y\,C_2\,F(-n_1+1,\gamma,y)+(\nu+\mu)\,C_1\,F(-n_1,\gamma,y)=0\,.$$

We apply the rule of differentiation of the confluent hypergeometric function

$$\frac{d}{dy}\,F(-n_1+1,\gamma,y)=\left(\frac{-n_1+1}{\gamma}-1\right)\,F(-n_1+1,\gamma+1,y)+F(-n_1+1,\gamma,y)\,,$$

and use the formula for contiguous confluent hypergeometric functions

$$y\,F(-n_1+1,\gamma+1,y)=\gamma F(-n_1+1,\gamma,y)-\gamma F(-n_1,\gamma,y)\,,$$

as a result we obtain

$$C_2\,\left(-n_1+1-\gamma\right)F(-n_1+1,\gamma,y)+C_2\,(A+\epsilon)\,F(-n_1+1,\gamma,y)-\\ -C_2\,\left(-n_1+1-\gamma\right)F(-n_1,\gamma,y)+(\nu+\mu)\,C_1\,F(-n_1,\gamma,y)=0\,.$$

Taking into account $-n_1=A-\epsilon$, we obtain

$$\frac{C_1}{C_2}=\frac{-n_1-2A}{\nu+\mu}=-\frac{A+\epsilon}{\nu+\mu}\,. \tag{71}$$

It is easily checked that two expressions for relative coefficients, (70) and (71), coincide. Indeed

$$\frac{C_1}{C_2}=-\frac{\nu-\mu}{A-\epsilon},\quad \frac{C_1}{C_2}=-\frac{A+\epsilon}{\nu+\mu}\;\Longrightarrow$$

$$\nu^2-\mu^2=A^2-\epsilon^2\quad\Longleftrightarrow\quad e^2\equiv e^2\,.$$

4 The spin $1/2$ particle in Coulomb field; solutions constructed completely by Heun functions

The idea to construct spectra within the Heun equation theory seems to be a very promising one. Let us try to consider, in this line of arguments, the known problem of a spin $1/2$ particle in the presence of external Coulomb field. To this end, let us turn again to the equations in presence of Coulomb potential (when $\delta = +1$)

$$\begin{aligned} \left(\frac{d}{dr} \, + \, \frac{\nu}{r}\right) f \, + \, \left(E + \frac{e}{r} \, + \, m\right) g \, = \, 0 \, , \\ \left(\frac{d}{dr} \, - \, \frac{\nu}{r}\right) g \, - \, \left(E + \frac{e}{r} \, - \, m\right) f \, = \, 0 \, . \end{aligned} \tag{72}$$

After eliminating the function g, one gets

$$\begin{aligned} \frac{d^2 f}{dr^2} + \frac{e}{r(Er + e + mr)} \frac{df}{dr} + \left[\frac{e(e^2 - \nu^2)}{r^2(Er + e + mr)}\right. \\ \left. + \frac{E\,(3e^2 - \nu^2) - \nu\,(m + E) + m\,(e^2 - \nu^2)}{r\,(Er + e + mr)}\right. \\ \left. + \frac{e\,(E + m)\,(3E - m)}{Er + e + mr} + \frac{r(E - m)(E + m)^2}{Er + e + mr}\right] f = 0 \, . \end{aligned} \tag{73}$$

After changing the variable

$$x = -\frac{(E + m)\, r}{e} \, , \tag{74}$$

equation (73) takes the form

$$x \, \frac{d^2 f}{dx^2} - \frac{1}{x - 1} \, \frac{df}{dx} + \left[\frac{e^2(Ex - mx - 2E)}{E + m} + \frac{e^2 - \nu^2}{x} - \frac{\nu}{x - 1}\right] f = 0 \, . \tag{75}$$

By separating the two factors $f(x) = x^A e^{Cx} F(x)$, one derives for F

$$\begin{aligned} \frac{d^2 F}{dx^2} + \left(2C + \frac{2A + 1}{x} - \frac{1}{x - 1}\right) \frac{dF}{dx} + \left[C^2 + \frac{e^2(E - m)}{E + m}\right. \\ \left. + \frac{A^2 + e^2 - \nu^2}{x^2} + \frac{A + C + 2AC - 2Ee^2/(E + m) + \nu}{x} - \frac{A + C + \nu}{x - 1}\right] F = 0 \, . \end{aligned} \tag{76}$$

When A, C are taken as (the bound states being of first interest)

$$\begin{aligned} C^2 + \frac{e^2(E - m)}{E + m} = 0 \qquad \Rightarrow \qquad C = +e\sqrt{\frac{m - E}{m + E}} \, , \\ A^2 + e^2 - \nu^2 = 0 \qquad \Rightarrow \qquad A = +\sqrt{\nu^2 - e^2} \, , \end{aligned} \tag{77}$$

equation (76) becomes simpler,

$$\begin{aligned} \frac{d^2 F}{dx^2} + \left(2C + \frac{2A + 1}{x} - \frac{1}{x - 1}\right) \frac{dF}{dx} \\ + \left[\frac{A + C + \nu + 2AC - 2Ee^2/(E + m)}{x} - \frac{A + C + \nu}{x - 1}\right] F = 0 \, , \end{aligned} \tag{78}$$

which is the confluent Heun equation for $F(\alpha,\beta,\gamma,\delta,\eta;x)$

$$\frac{d^2}{dz^2}F+\left(a+\frac{\beta+1}{z}+\frac{\gamma+1}{z-1}\right)\frac{dF}{dz}$$
$$+\left(\frac{1}{2}\frac{a+a\gamma+\beta+\beta\gamma+\gamma+2\delta+2\eta}{z-1}+\frac{1}{2}\frac{a\beta+a-\beta\gamma-\beta-\gamma-2\eta}{z}\right)F=0 \tag{79}$$

with parameters determined by

$$a=2C=+2e\sqrt{\frac{m-E}{m+E}}\,,\qquad \beta=2A=+2\sqrt{\nu^2-e^2}\,,$$
$$\gamma=-2\,,\qquad \delta=-\frac{2Ee^2}{E+m}\,,\qquad \eta=1-\nu+\frac{2Ee^2}{E+m}\,. \tag{80}$$

The known condition to obtain polynomials is

$$\delta=-a\left(n+\frac{\gamma+\beta+2}{2}\right)\,. \tag{81}$$

which can bee written as

$$-\frac{2Ee^2}{E+m}=-2e\sqrt{\frac{m-E}{m+E}}\,(\,n+\sqrt{\nu^2-e^2}\,)\,, \tag{82}$$

or

$$\frac{Ee}{\sqrt{m^2-E^2}}=N\,,\qquad N=n+\sqrt{\nu^2-e^2}\,; \tag{83}$$

from where it follows

$$E=\frac{m}{\sqrt{1+e^2/N^2}}\,. \tag{84}$$

This is the exact energy spectrum for the hydrogen atom in Dirac theory.

5 Conclusions

In the paper, the well-known quantum mechanical problem of a spin 1/2 particle in external Coulomb potential, reduced to a system of two first-order differential equations, is studied from the point of view of possible applications of the Heun function theory to treat this system. It is shown that in addition to the standard way to solve the problem in terms of the confluent hypergeometric functions (proposed in 1928 by G. Darvin and W. Gordon), there are possible several other possibilities which rely on applying the confluent Heun functions. Namely, in the paper there are elaborated two combined possibilities to construct solutions: the first applies when one equation of the pair of relevant functions is expressed trough hypergeometric functions, and another constructed in terms of confluent Heun functions. In this respect, certain relations between the two classes of functions are established. It is shown that both functions of the system may be expressed in terms of confluent Heun functions. All the ways to study this problem lead us to a single energy spectrum, which indicates their correctness.

References

[1] K. Heun. Zur Theorie der Riemann'schen Functionen Zweiter Ordnung mit Verzweigungspunkten. Math. Ann. **33**. 161-179 (1889).

[2] A. Ronveaux (ed.). Heun's differential equation. Oxford University Press, Oxford (1995).

[3] P.S. Epstein. The stark effect from the point of view of Schrdinger quantum theory. Phys. Rev. **28**. 695 (1926).

[4] W.G. Baber, H.R. Hassé. The two centre problem in wave mechanics. Proc. Cambridge Philos. Soc. **31**. 564–581 (1935).

[5] M.F. Manning. Exact solutions of the Schrödinger equation. Phys. Rev. **48**. 161–164 (1935).

[6] T. Regge, J.A. Wheeler. Stability of a Schwarzschild Singularity. Phys. Rev. **108**. 1063–1069 (1957).

[7] A. Lamieux and A.K. Bose. Construction de potentiels pour lesquelles léquation de Schrödinger est soluble. Ann. Inst. Henri Poincaré. **A 10**. 259–270 (1969).

[8] S.V. Dhurandhar, C.V. Vishveshwara and J.M. Cohen. Electromagnetic, neutrino and gravitational fields in the Kasner space-time with rotational symmetry. Class. Quantum Grav. **1**. 61–69 (1984).

[9] R.N. Chaudhuri and S. Mukherjee. On the $\mu x^2 + \lambda x^4 + \eta x^6$ interaction. J. Phys. **A 17**. 3327–(1984).

[10] B. Léauté, G. Marcilhacy. On the Schrödinger equations of rotating harmonic, threedimensional and doubly anharmonic oscillators and a class of confinement potentials in connection with the biconfluent Heun differential equation. J. Phys. **A 19**. 3527–3533 (1986).

[11] Leaver E L. Solutions to a generalized spheroidal wave equation: Teukolsky equations in general relativity, and the two-center problem in molecular quantum mechanics. J. Math. Phys. **27**. 1238–1265 (1986).

[12] R. Pons, G. Marcilhacy. Exact solutions for electromagnetic, neutrino and gravitational fields in Kasner spacetime. Class. Quantum Grav. **4**. 171-179 (1987).

[13] I. Costa, N. Derruelle, M. Novello, N. F. Svaiter. Quantum fields in cosmological spacetimes: a soluble example. Class. Quantum Grav. **6**. 1893–1907 (1989).

[14] P.P. Ray, K. Mahata. Bounded states of the potential $V(r) = \frac{Ze^2}{r+\beta}$. J. Phys. A: Math. Gen. **22**. 3161 (1989).

[15] H. Exton. The exact solution of two new types of Schrodinger equation. J. Phys. **A 28**. 6739–6741 (1995).

[16] V.S. Otchik. Analytic solutions of the Teukolsky equations in Quantum Systems: New Trends and Methods, edited by A. O. Barut et al. (World Scientific, Singapore, 1995).

[17] H. Suzuki, E. Takasugi, H. Umetsu. Perturbations of Kerr-de Sitter black holes and Heun's equations. Progr. Theoret. Phys. **100**. 491–505 (1998); arXiv:gr-qc/9805064.

[18] T.T. Truong and D. Bazzali. Exact low-lying states of two interacting equally charged particles in a magnetic field. Phys. Lett. **A 269**. 186 (2000).

[19] G. Esposito, R. Roychowdhury. On the complete analytic structure of the massive gravitino propagator in four-dimensional de Sitter space. Gen. Rel. Grav. **42**. 1221–1238 (2010); arXiv:0907.3634 [hep-th].

[20] A.M. Ishkhanyan, K.-A. Suominen. Solutions of the two-level problem in terms of biconfluent Heun functions. J. Phys. **A 34**. 6301–6306 (2001).

[21] A.M. Ishkhanyan, K.-A. Suominen. Analytic treatment of the polariton problem for a smooth interface. J. Phys. A: Math. Gen. **34**. L591-L598 (2001).

[22] A. Ralko, T.T.Truong. Heun Functions and the energy spectrum of a charged particle on a sphere under magnetic field and Coulomb force. J. Phys. A: Math. Gen. **35**. 9573-9584 (2002); arXiv:quant-ph/0209152

[23] S.Q. Wu, X. Cai. Massive complex scalar field in a Kerr-Sen Black Hole Background: exact solution of waveequation and Hawking radiation. J. Math. Phys. **44**. 1084-1988 (2003); arXiv:gr-qc/0303075

[24] T. Birkandan and M. Hortacsu. Heun and Mathieu functions as solutions of the Dirac equation. EAS Publ.Ser. **30**. 265 (2008); arXiv:0710.4243 [gr-qc].

[25] Takeshi Oota, Yukinori Yasui. Toric Sasaki-Einstein manifolds and Heun equation. Nucl.Phys. **B 742**. 275–294 (2006); arXiv:hep-th/0512124.

[26] P.P. Fiziev. Exact solutions of Regge-Wheeler equation and quasi-normal modes of compact objects. Class. Quantum Grav. **23**. 2447–2468 (2006).

[27] T. Birkandan, M. Hortacsu. Examples of Heun and Mathieu Functions as Solutions of Wave Equations in Curved Spaces. J.Phys. **A40**. 1105–1116 (2007); arXiv:gr-qc/0607108v3.

[28] David Petroff. Slowly Rotating Homogeneous Stars and the Heun Equation. Class. Quant. Grav. **24**. 1055–1068 (2007); arXiv:gr-qc/0701081v2 6 Feb 2007

[29] P.P. Fiziev. Exact Solutions of Regge-Wheeler Equation. J. Phys. Conf. Ser. **66**. 012016 (2007); arXiv:gr-qc/0702014v1.

[30] D. Bouaziz, M. Bawin. Regularization of the singular inverse square potential in quantum mechanics with a Minimal length. Phys. Rev. **A76**. 032112 (2007), arXiv:0711.0599 [quant-ph].

[31] T. Takemura. Finite-gap potential, Hein differential equation and WKB analysis. arXiv:math/0703256v1 [math.CA] 9 Mar 2007

[32] R. Giachetti, E. Sorace. States of the Dirac equation in confining potentials. Phys. Rev. Lett. **101**. 190401 (2008); arXiv:0706.0127 [hep-th].

[33] A.Al-Badawi, I.Sakalli. Solution of the Dirac equation in the rotating Bertotti-Robinson spacetime. J. Math. Phys. **49**. 052501 (2008); arXiv:0805.4485 [gr-qc].

[34] Lea Jaccoud El-Jaick, Bartolomeu D. B. Figueiredo. A limit of the confluent Heun equation and the Schrödinger equation for an inverted potential and for an electric dipole. J. Math. Phys. **50**. 123511 (2009).

[35] A.M. Ishkhanyan, R. Sokhoyan, B. Joulakian, and K.-A. Suominen: Rosen-Zener model in cold molecule formation? Opt. Commun. **282**. 218–226 (2009).

[36] A. Ishkhanyan, R. Sokhoyan, K.-A. Suominen, C. Leroy, and H.-R. Jauslin: Quadratic-nonlinear Landau-Zener transition for association of an atomic Bose-Einstein condensate with inter-particle elastic interactions included Eur. Phys. J. **D 56**. 421–429 (2010).

[37] A. Arda, O. Aydogdu, R. Sever. Scattering of Woods-Saxon potential in Schrodinger equation. J. Phs. A: Math Theor. **43**, 425204 (2010); arXiv:1009.5181 [quant-ph]

[38] Javier Sesma, The generalized quantum isotonic oscillator. J. Phys. A: Math. Theor. **43**. 185303 (2010); arXiv:1005.1227 [quant-ph]

[39] Richard L. Hall, Nasser Saad, K.D. Sen, Soft-core Coulomb Potential and Heun's Differential Equation. J. Math. Phys. **51** (2010) 022107. arXiv:0912.3445

[40] Roumen S. Borissov, Plamen P. Fiziev, Exact solutions of Teukolsky master equation with continuous spectrum. Bulg. J. Phys. **37**. 065–089 (2010); arXiv:0903.3617 [gr-qc].

[41] Fiziev, P.P.: Classes of exact solutions to the Teukolsky master equations. Class. Quantum Grav. **27**. 135001 (2010); arXiv:gr-qc/0908.4234.

[42] V.V. Kisel, E.M. Ovsiyuk, V.M. Red'kov. On the wave functions and energy spectrum for a spin 1 particle in external Coulomb field. NPCS. **13**. 352–367 (2010.)

[43] V.H. Zaveri. Quarkonium and hydrogen spectra with spin-dependent relativistic wave equation. Pramana J. Phys. **75**. 579–598 (2010).

[44] P. Fiziev, D. Staicova. Application of the confluent Heun functions for finding the quasinormal modes of nonrotating black holes. Phys. Rev. **D 84**. 127502 (2011); arXiv:1109.1532 [gr-qc]

[45] M.S. Cunha, H.R. Christiansen. Confluent Heun functions in gauge theories on thick braneworlds. Phys. Rev. **D84**. 085002 (2011); arXiv:1109.3486 [hep-th].

[46] V.V. Kisel, G.G. Krylov, E.M. Ovsiyuk, M. Amirfakchrian, V.M. Red'kov. Wave functions of the particle with polarizability in Coulomb field. Nonlinear Dynamics and Applications. Vol. 18, P. 168–179, Minsk, 2011; Proc. of XVIII Annual Seminar NPCS-2011, May 17–20, 2011, Minsk, Belarus; 13 pages.

[47] E. Ovsiyuk, O. Veko and M. Amirfakchrian. On Schrodinger equation with potential $U = -\alpha r^{-1} + \beta r + kr^2$ and the biconfluent Heun functions theory. Nonlinear Phenomena in Complex Systems. **15**, 163 (2012).

[48] E.M. Ovsiyuk. On solutions of Maxwell equations in the space-time of Schwarzschild black hole. NPCS. **15**. 81–91 (2012).

[49] E.M. Ovsiyuk, O.V. Veko, V.V. Kisel, V.M. Red'kov. New problems of quantum mechanics and Heun equation. Naucno-tekchnicheskie vedomosti St.Petersburg University. Ser. Fiz.-mat. **1(141)**. 137–145 (2012).

[50] E.Ovsiyuk, O.Veko, K.Kazmerchuk, V.Kisel, V.Red'kov Quantum mechanics of a spin 1 particle in the magnetic monopole potential, in spaces of Euclid, Lobachevsky, and Riemann: nonrelativistic approximation. Ukr. Phys. J. **58**. 1073–1083 (2013).

[51] T.A. Shahverdyan, D.S. Mogilevtsev, A.M. Ishkhanyan, V.M. Red'kov A.M. Complete-return spectrum for a generalized Rosen-Zener two-state term-crossing model. Nonlinear Phenomena in Complex Systems. **16**. 86–92 (2013)

[52] V.M. Red'kov, A.V. Chichurin. A Symbolic-Numerical Method for Solving the Differential Equation Describing the States of Polarizable Particle in Coulomb Potential Programming and Computer Software, 2014, Vol. 40, no. 2, P. 86–92. Pleiades Publishing, Ltd., 2014.

[53] A. Chychuryn, V. Red'kov. Quantum mechanical scalar particle with polarisability in the Coulomb field, analytical and numerical consideration. Studia i Materialy. 2013. Vol. 6. S. 73–89. Europejska Wyzsza Szkola Informtyczno-Ekonomiczna, Warszawa.

[54] A.A. Bogush, G.G. Krylov, E.M. Ovsiyuk, and V.M. Red'kov. Maxwell equations in complex form of Majorana-Oppenheimer, solutions with cylindric symmetry in Riemann S_3 and Lobachevsky H_3 spaces. Ricerche di matematica. **59**. 59–96 (2010).

[55] V.M. Red'kov, E.M. Ovsiyuk. On exact solutions for quantum particles with spin S = 0, 1/2, 1 and de Sitter event horizon. Ricerche di matematica. **60**. 57–88 (2011).

[56] Red'kov, E. Ovsiyuk, O. Veko. Spin 1/2 particle in the field of Dirac string on the background of de Sitter space–time. Uzhhorod University Scientific Herald. Series Physics. **32**. 141–150 (2012).

[57] E.M. Ovsiyuk, O.V. Veko, V.M. Red'kov. On simulating a medium with special reflecting properties by Lobachevsky geometry. NPCS. (2013). **16**. 331–344

[58] E. Ovsiyuk, O. Florea, A. Chichurin, V. Red'kov Electromagnetic Field in Schwarzschild Black Hole Background. Analytical Treatment and Numerical Simulation. Computer Algebra Systems in Teaching and Research, Vol. IV, No 1, Siedlce, Publ. Collegium Mazovia, 2013, P. 204 -214. [Proceedings of the 7-th International Workshop on Computer Algebra Systems in Teaching and Research (CASTR-2113). Siedlce, Poland, September 22–25, 2013.]

[59] E.R. Figueiredo Medeiros, E.R. Bezerra de Mello. Relativistic quantum dynamics of a charged particle in cosmic string spacetime in the presence of magnetic field and scalar potential. Eur. Phys. J. **C 72**. 2051 (2012); arXiv:1108.3786 [hep-th]

[60] M. Hortacsu. Heun Functions and their uses in Physics. pp. 23–39 in: Proceedings of the 13th Regional Conference on Mathematical Physics, Antalya, Turkey, October 27-31, 2010, Edited by Ugur Camci and Ibrahim Semiz, pp. 23-39. World Scientific (2013); arXiv:1101.0471 [math-ph].

[61] T. Birkandan, M. Hortacsu. Examples of Heun and Mathieu Functions as Solutions of Wave Equations in Curved Spaces. J. Phys. A: Math. Theor. **40**. 1105–1116 (2007); arxiv: 0607108 [gr-qc]

[62] D. Momeni, K. Yerzhanov, R. Myrzakulov, Quantized Black Hole and Heun function. Can. J. Phys. 90: 877-881 (2012), arXiv:1009.0130 [physics.gen-ph]

[63] C. A. Downing. On a solution of the Schrodinger equation with a hyperbolic double-well potential. J. Math. Phys. **54**. 072101 (2013); arXiv:1211.0913,

[64] D. Batic, R. Williams, M. Nowakowski. Potentials of the Heun class. J. Phys. **A 46**. 245204 (2013).

[65] A.A. Rajabi, M. Hamzavi. Relativistic effect of external magnetic and Aharonov-Bohm fields on the unequal scalar and vector Cornell model. Eur. Phys. J. Plus **128**. 5 (2013).

[66] M. Hamzavi, A.A. Rajabi. Scalar-vector-pseudoscalar Cornell potential for a spin-1/2 particle under spin and pseudospin symmetries: 1+1 dimensions. Ann. Phys. **334**. 316 (2013).

[67] V.I. Osherov, V.G. Ushakov, Stark problem in terms of the Stokes multipliers for the triconfluent Heun equation. Phys. Rev. **A 88**. 053414 (2013).

[68] H. S. Vieira, V. B. Bezerra, C. R. Muniz. Exact solutions of the Klein-Gordon equation in the Kerr-Newman background and Hawking radiation. Annals of Physics. **350**. 14–28 (2014).

[69] F. Caruso, J. Martins, V. Oguri. Solving a two-electron quantum dot model in terms of polynomial solutions of a Biconfluent Heun equation. Annals of Physics. **347**. 130 (2014).

[70] J. Karwoswki and H.A. Witek. Biconfluent Heun equation in quantum chemistry: Harmonium and related systems. Theor. Chem. Accounts. **133**. Paper 1494 [11 pages] (2014).

[71] J. M. Hoff da Silva, S. H. Pereira. Exact solutions to Elko spinors in spatially flat Friedmann-Robertson-Walker spacetimes. Journal of Cosmology and Astroparticle Physics. **3**. 009 (2014); arXiv:1401.3252 [hep-th].

[72] O.V. Veko, K.V. Kazmerchuk, E.M. Ovsiyuk, V.M. Red'kov, A.M. Ishkhanyan. Hydrogen atom in de Sitter spaces and Heun functions. Report to International Conference and Workshop "Quanta and Matter: Through Physics to Future Emerging Technologies 22-26 September 2014. Yerevan-Tsaghkadzor, Armenia.

[73] Berestetzkiy, V.B.; Lifshitz, E.M.; Pitaevskiy, L.B., Quantum Electrodynamics. Moscow, Nauka, 1989 (in Russian)

[74] V.M. Red'kov, E.M. Ovsiyuk. Quantum mechanics in spaces of constant curvature. Nova Science Publishers. Inc., New York, 2012.

[75] Ovsiyuk,E.M.; Kisel, V.V.; Red'kov, V.M.: Maxwell Electrodynamics and Boson Fields in Spaces of Constant Curvature. Nova Science Publishers Inc., New York, 2014. 486 pages.

[76] Varshalovich, D.A., Moskalev, A.N., Hersonskiy, V.K.: Quantum theory of angular moment. Nauka, Leningrad, 1975 (in Russian).